

UT-08-17

IPMU-08-0029

A Measurement of Neutralino Mass at the LHC in Light Gravitino Scenarios

Koichi Hamaguchi^{1,2}, Eita Nakamura¹ and Satoshi Shirai¹

¹ *Department of Physics, University of Tokyo,
Tokyo 113-0033, Japan*

² *Institute for the Physics and Mathematics of the Universe, University of Tokyo,
Chiba 277-8568, Japan*

Abstract

We consider supersymmetric (SUSY) models in which a very light gravitino is the lightest SUSY particle. Assuming that a neutralino is the next-to-lightest SUSY particle, we present a measurement of the neutralino mass at the LHC in two photons + missing energy events, which is based on the M_{T2} method. It is a direct measurement of the neutralino mass itself, independent of other SUSY particle masses and patterns of cascade decays before the neutralino is produced.

Among various supersymmetric (SUSY) models, those with an ultralight gravitino of mass $m_{3/2} \lesssim \mathcal{O}(10)$ eV are very attractive, since they are completely free from notorious gravitino problems [1]. In this letter, we assume a neutralino is the next-to-lightest SUSY particle (NLSP), and present a measurement of its mass at the LHC. It is based on the so-called M_{T2} method [2]. We show that this method can directly determine the neutralino mass, independently of other SUSY particle masses, and it does not rely on specific patterns of cascade decays before the neutralino is produced.

In the scenario considered here, essentially all the SUSY events will end up with two neutralino NLSPs,¹ each of which then dominantly decays into a gravitino and a photon.² We assume that the decay length of the NLSP neutralino is so short that the decay occurs inside the detector and the photons' momenta are measured well. Therefore, the main signature at the LHC will be two high transverse momentum photons and a large missing transverse momentum. If such a signal will indeed be discovered, one of the most natural candidates for the underlying model is a SUSY model with a gravitino LSP and a neutralino NLSP.

Furthermore, from the prompt decay of the neutralino, we can assume that the gravitino is very light, essentially massless for the following discussion. This is because the NLSP decay length is proportional to the gravitino mass squared as

$$c\tau_{\text{NLSP}} \sim 20\mu\text{m} \left(\frac{m_{3/2}}{1\text{ eV}}\right)^2 \left(\frac{m_{\text{NLSP}}}{100\text{ GeV}}\right)^{-5}, \quad (1)$$

and a heavier gravitino ($m_{3/2} > \mathcal{O}(1)$ keV) would make the neutralino decay outside the detector.³ This indirect information of the massless LSP plays a crucial role in the NLSP mass determination.

Let us start by briefly explaining the M_{T2} method [2]. Suppose that there is a particle A which promptly decays by the process $A \rightarrow B + X$, where B is a visible (standard-model) particle and X is a neutral and undetected particle. When two A s are produced in a collider, we can measure the two B s' transverse momenta $\mathbf{p}_T^{B,1}$, $\mathbf{p}_T^{B,2}$ and the missing

¹We assume R -parity conservation.

²We do not discuss the case in which the neutralino mainly decays into a Higgs/ Z -boson and a gravitino.

³For a moderate gravitino mass corresponding $c\tau_{\text{NLSP}} = \mathcal{O}(10)$ cm $- \mathcal{O}(10)$ m, the neutralino decay causes “non-pointing” photons [3]. The present method of the neutralino mass determination may also work in this case.

transverse momentum $\mathbf{p}_T^{\text{miss}} = \mathbf{p}_T^{X,1} + \mathbf{p}_T^{X,2}$.⁴ The M_{T2} variable is then given by

$$(M_{T2})^2 \equiv \min_{\mathbf{p}_T^{\text{miss},1} + \mathbf{p}_T^{\text{miss},2} = \mathbf{p}_T^{\text{miss}}} \left[\max \left\{ (M_T^{(1)})^2, (M_T^{(2)})^2 \right\} \right], \quad (2)$$

where the minimization is taken over all possible momentum splittings, and

$$(M_T^{(i)})^2 = m_B^2 + m_X^2 + 2(E_T^{\text{miss},i} E_T^{B,i} - \mathbf{p}_T^{\text{miss},i} \cdot \mathbf{p}_T^{B,i}) \quad \text{for } i = 1, 2 \quad (3)$$

with $E_T^{\text{miss},i} \equiv \sqrt{m_X^2 + |\mathbf{p}_T^{\text{miss},i}|^2}$ and $E_T^{B,i} \equiv \sqrt{m_B^2 + |\mathbf{p}_T^{B,i}|^2}$. This M_{T2} variable is designed to have the endpoint at m_A when we input the correct value of m_X . However, in general, the mass m_X of the missing particle X is unknown, and therefore one can obtain only a relation between m_X and m_A .⁵

A crucial point in the scenario considered here ($\{A, B, X\} = \{\text{neutralino}, \gamma, \text{gravitino}\}$) is that we can assume the massless LSP, $m_X = m_{3/2} = 0$, as discussed above. Therefore, we can directly determine the NLSP mass by the M_{T2} method. As we show in the Appendix, the M_{T2} variable in this case is analytically expressed as⁶

$$(M_{T2})^2 = \begin{cases} 2p_1 p_2 z & \text{for } c_1 < 0 \text{ or } c_2 < 0 \\ 0 & \text{for } c_1 \geq 0 \text{ and } c_2 \geq 0 \end{cases}, \quad (4)$$

where $p_1 \equiv |\mathbf{p}_T^{\gamma,1}|$, $p_2 \equiv |\mathbf{p}_T^{\gamma,2}|$, c_1 and c_2 are given by

$$\mathbf{p}_T^{\text{miss}} = c_1 \mathbf{p}_T^{\gamma,1} + c_2 \mathbf{p}_T^{\gamma,2}, \quad (5)$$

and z is a real positive solution of the following equations:

$$\begin{aligned} 4(a-b)^2 &= (\sqrt{2(a+b)+3}-1)(\sqrt{2(a+b)+3}+3)^3, \\ a &= \frac{1}{r^{\frac{1}{3}}} \left[\frac{r}{2} - \cos \theta + c_1 \sin^2 \theta \frac{1}{z} \right], \\ b &= r^{\frac{1}{3}} \left[\frac{1}{2r} - \cos \theta + c_2 \sin^2 \theta \frac{1}{z} \right], \\ r &= \frac{p_2}{p_1}, \quad \cos \theta = \frac{\mathbf{p}_T^{\gamma,1} \cdot \mathbf{p}_T^{\gamma,2}}{p_1 p_2}. \end{aligned} \quad (6)$$

⁴We assume that the missing $\mathbf{p}_T$ is dominantly caused by the two X s and the contribution of other sources of missing $\mathbf{p}_T$ are negligible.

⁵See also recent developments in the M_{T2} method [4].

⁶For completeness, we also show an analytic expression of M_{T2} for the case of massive LSP ($m_X \neq 0$) in the Appendix.




Figure 1: Mass spectrum of SIGM

Note that M_{T2} is completely defined by the missing transverse momentum and photon momenta, independently of other kinematical variables. We should also emphasize that the present method does not rely on a direct pair-production of the NLSPs, i.e., we do not assume back-to-back transverse momenta of the NLSPs, $\mathbf{p}_T^{\text{miss}} + \mathbf{p}_T^{\gamma,1} + \mathbf{p}_T^{\gamma,2} = 0$. In the following, we show how this method works at the LHC, by taking explicit examples of gauge mediated SUSY breaking (GMSB) models which realize the mass spectrum with an ultralight gravitino LSP and a neutralino NLSP.

We consider two gauge mediation models for a demonstration. In the following, mass spectrums are calculated by ISAJET 7.72 [5] and we use programs Herwig 6.5 [6] and AcerDET-1.0 [7] to simulate LHC signatures.

The first example is a strongly interacting gauge mediation (SIGM) model [8], in which the NLSP is a neutralino. We take the same SIGM parameters as the example in Sec. 4 of Ref. [8]. The mass spectrum is shown in Fig. 1. The masses of the lightest neutralino and gravitino are 356 GeV and 10 eV, respectively.

We take the events cuts as follows:

- ≥ 4 jets with $p_T > 50$ GeV and $p_{T,1,2} > 100$ GeV.
- ≥ 2 photons with $p_T > 20$ GeV.




Figure 2: A distribution of M_{T2} for the SIGM example. (a): parton level signature. (b): detector level signature.

- $M_{\text{eff}} > 500$ GeV, where

$$M_{\text{eff}} = \sum_{\text{jets}}^4 p_{Tj} + p_T^{\text{miss}}. \quad (7)$$

- $p_T^{\text{miss}} > 0.2M_{\text{eff}}$.

Under these cuts, we see that the standard-model backgrounds are almost negligible.

In Fig. 2-(a), a parton level distribution of M_{T2} is shown for an integrated luminosity of 10 fb^{-1} . Here, we take the sum of gravitino and neutrino transverse momenta as the parton level missing $\mathbf{p}_T$. As discussed in Ref. [8], very little number of leptons are produced in the SIGM. Therefore, missing $\mathbf{p}_T$ is due to almost only gravitinos and the assumption that $\mathbf{p}_T^{\text{miss}} = \mathbf{p}_T^{\text{LSP1}} + \mathbf{p}_T^{\text{LSP2}}$ is satisfied. There is a clear edge at $M_{T2} \simeq m_{\tilde{\chi}_1^0} = 356$ GeV.

In Fig. 2-(b), we show a distribution of M_{T2} after taking account of detector effects. In order to extract the point of the edge, we use a simple fitting function;

$$f(x) = (ax + b)\theta(-x + M) + (cx + d)\theta(x - M), \quad (8)$$

where $\theta(x)$ is the step function and a, b, c, d and M are fitting parameters. We fit the data with $f(x)$ over $300 \leq M_{T2} \leq 500$ GeV and find

$$m_{\tilde{\chi}_1^0} = 357 \pm 3 \text{ GeV}. \quad (9)$$




Figure 3: Mass spectrum of SPS8

Here, the estimation of the error is done by ‘eye’ because of lack of information on the shape of the M_{T2} distribution. The estimation that $m_{\tilde{\chi}_1^0} = 357 \pm 3$ GeV is very good agreement with the true value $m_{\tilde{\chi}_1^0} = 356$ GeV.

Next we show another example. We study the Snowmass benchmark point SPS8 [9], which is a minimal gauge mediation model with a neutralino NLSP. In Fig. 3, SPS8 mass spectrum is shown. The masses of the lightest neutralino and gravitino are 139 GeV and 4.8 eV, respectively.

In Fig. 4(a), a parton level distribution of M_{T2} is shown for an integrated luminosity of 10 fb^{-1} . The event cuts are the same as in the previous SIGM case. The blue and dashed line represents the case that $\mathbf{p}_T^{\text{miss}} = \sum_{\text{gravitino}} \mathbf{p}_T$ and the red and solid line $\mathbf{p}_T^{\text{miss}} = \sum_{\text{gravitino}} \mathbf{p}_T + \sum_{\text{neutrino}} \mathbf{p}_T$. In SPS8, there are many neutrino production sources. Hence, we cannot see a clear edge as in the SIGM case. However, there is a cliff at $M_{T2} \simeq m_{\tilde{\chi}_1^0} = 139$ GeV.

In Fig. 4(b), detector level distribution of M_{T2} is shown. To get the value of $m_{\tilde{\chi}_1^0}$, we fit the data with $f(x)$ in Eq. (8) over $110 \leq M_{T2} \leq 180$ GeV. Then we get

$$m_{\tilde{\chi}_1^0} = 139 \pm 3 \text{ GeV}. \quad (10)$$




Figure 4: A distribution of M_{T2} for the SPS8. (a): parton level signature. The blue and dashed line represents the case that $\mathbf{p}_T^{\text{miss}} = \sum_{\text{gravitino}} \mathbf{p}_T$ and the red and solid line $\mathbf{p}_T^{\text{miss}} = \sum_{\text{gravitino}} \mathbf{p}_T + \sum_{\text{neutrino}} \mathbf{p}_T$. (b): detector level signature.

The error estimation is done by ‘eye’. This value agrees with the true value ($m_{\tilde{\chi}_1^0} = 139$ GeV).

In summary, we have presented a determination of the neutralino mass for the SUSY models with an ultralight gravitino LSP and a neutralino NLSP, which may work in the early stage of the LHC.

Though we have considered GMSB models with a neutralino NLSP, our method is applicable to any model in which the signal events will lead to a pair of cascade decays that result in

$$\cdots \rightarrow \text{any cascade decay} \rightarrow A \rightarrow B + X, \quad (11)$$

where B is a visible (standard-model) particle and X is a missing particle that is almost massless. The mass of A is then determined by the two B s’ momenta and the missing transverse momentum.

For example, let us consider GMSB models with a slepton NLSP. In this case, the slepton, lepton and gravitino correspond to A , B and X in Eq. (11), respectively. In addition to leptons from the sleptons’ decays, many other leptons are produced in this scenario. However, we may see which of observed leptons is produced through the slepton decay by measuring lepton’s momentum, or by detecting a kink of its track for a long-

lived slepton. In such a case, we can measure the slepton mass with the M_{T2} method as discussed above. Furthermore, the present method may work in an axino LSP scenario.

Acknowledgements

We thank Tsutomu Yanagida for useful discussion. This work was supported by World Premier International Center Initiative (WPI Program), MEXT, Japan. The work by KH is supported by JSPS (18840012). The work of SS is supported in part by JSPS Research Fellowships for Young Scientists.

Appendix

In this appendix we derive Eq. (4). We start from Eqs. (2) and (3). We assume that B is massless.

(i) $m_X = 0$ case: First, we consider the case that X is a massless particle. The M_{T2} variable is defined by Eqs. (2) and (3) with $m_B = m_X = 0$. If a momentum splitting is the correct one, i.e., $\mathbf{p}_T^{\text{miss},1} = \mathbf{p}_T^{X,1}$ and $\mathbf{p}_T^{\text{miss},2} = \mathbf{p}_T^{X,2}$, then each transverse mass is smaller than the mass of A , m_A :

$$m_A^2 = (p^{B,i} + p^{X,i})^2 = 2(|\mathbf{p}_T^{X,i}| |\mathbf{p}_T^{B,i}| \cosh(\Delta y^i) - \mathbf{p}_T^{X,i} \cdot \mathbf{p}_T^{B,i}) \geq (M_T^{(i)})^2 \quad (12)$$

for $i = 1, 2$, where Δy^i is the rapidity difference of B and X in each decay chain. From this, it is clear that

$$M_{T2} \leq m_A. \quad (13)$$

We do not assume the relation

$$\mathbf{p}_T^{B,1} + \mathbf{p}_T^{B,2} = -\mathbf{p}_T^{\text{miss}}, \quad (14)$$

which holds in the case of a “back-to-back” pair production of A s. We may assume that $\mathbf{p}_T^{B,1}$ and $\mathbf{p}_T^{B,2}$ are linearly independent and $\mathbf{p}_T^{\text{miss}}$ can be expressed as

$$\mathbf{p}_T^{\text{miss}} = c_1 \mathbf{p}_T^{B,1} + c_2 \mathbf{p}_T^{B,2}. \quad (15)$$

Here, c_1 and c_2 are real coefficients and they are given by

$$c_1 = \frac{1}{\sin^2 \theta} \left[\frac{\mathbf{p}_T^{\text{miss}} \cdot \mathbf{p}_T^{B,1}}{(p_1)^2} - \frac{\mathbf{p}_T^{\text{miss}} \cdot \mathbf{p}_T^{B,2}}{p_1 p_2} \cos \theta \right], \quad (16)$$

$$c_2 = \frac{1}{\sin^2 \theta} \left[\frac{\mathbf{p}_T^{\text{miss}} \cdot \mathbf{p}_T^{B,2}}{(p_2)^2} - \frac{\mathbf{p}_T^{\text{miss}} \cdot \mathbf{p}_T^{B,1}}{p_1 p_2} \cos \theta \right], \quad (17)$$

where

$$p_1 \equiv |\mathbf{p}_T^{B,1}|, \quad p_2 \equiv |\mathbf{p}_T^{B,2}|, \quad \cos \theta \equiv \frac{\mathbf{p}_T^{B,1} \cdot \mathbf{p}_T^{B,2}}{p_1 p_2}. \quad (18)$$

The momentum splitting $\mathbf{p}_T^{\text{miss},1}$ and $\mathbf{p}_T^{\text{miss},2}$ can also be expressed as

$$\mathbf{p}_T^{\text{miss},1} = (c_1 - x)\mathbf{p}_T^{B,1} + y\mathbf{p}_T^{B,2}, \quad (19)$$

$$\mathbf{p}_T^{\text{miss},2} = x\mathbf{p}_T^{B,1} + (c_2 - y)\mathbf{p}_T^{B,2}, \quad (20)$$

where x and y are real variables. We rewrite Eq. (2) as

$$(M_{T2})^2 = 2p_1 p_2 \min_{x,y \in \mathbf{R}} \left[\max \left\{ z_1(x, y), z_2(x, y) \right\} \right], \quad (21)$$

where

$$z_1(x, y) \equiv \frac{(M_T^{(1)}(x, y))^2}{2p_1 p_2} = \sqrt{\left[\frac{c_1 - x}{r} + y \cos \theta \right]^2 + y^2 \sin^2 \theta} - \left[\frac{c_1 - x}{r} + y \cos \theta \right], \quad (22)$$

$$z_2(x, y) \equiv \frac{(M_T^{(2)}(x, y))^2}{2p_1 p_2} = \sqrt{[x \cos \theta + (c_2 - y)r]^2 + x^2 \sin^2 \theta} - [x \cos \theta + (c_2 - y)r], \quad (23)$$

and $r \equiv p_2/p_1$. It is clear that

$$z_1(x, y) \geq 0, \quad \text{and} \quad z_1(x, y) = 0 \Leftrightarrow y = 0 \ \& \ x \leq c_1, \quad (24)$$

$$z_2(x, y) \geq 0, \quad \text{and} \quad z_2(x, y) = 0 \Leftrightarrow x = 0 \ \& \ y \leq c_2. \quad (25)$$

From this, we can infer that

$$(M_{T2})^2 = 0 \quad \text{if} \quad c_1 \geq 0 \ \& \ c_2 \geq 0, \quad (26)$$

and for other values of c_1 and c_2 , $(M_{T2})^2$ is given by $(M_T^{(1)}(x, y))^2 = (M_T^{(2)}(x, y))^2$ at the point $(x, y) = (x_0, y_0)$ where the contours of $z_1(x, y)$ and $z_2(x, y)$ in the x - y plane become tangent to each other. We denote the corresponding value $z \equiv z_1(x_0, y_0) = z_2(x_0, y_0)$ in the following. Eqs. (22) and (23) yield

$$x_0 = -\frac{r \sin^2 \theta}{2z} \left(y_0 - \frac{z \cos \theta}{\sin^2 \theta} \right)^2 + \frac{rz}{2 \sin^2 \theta} + c_1, \quad (27)$$

$$y_0 = -\frac{\sin^2 \theta}{2rz} \left(x_0 - \frac{z \cos \theta}{\sin^2 \theta} \right)^2 + \frac{z}{2r \sin^2 \theta} + c_2 \quad (28)$$

with the tangential condition

$$\frac{\sin^4 \theta}{z^2} \left(x_0 - \frac{z \cos \theta}{\sin^2 \theta} \right) \left(y_0 - \frac{z \cos \theta}{\sin^2 \theta} \right) = 1. \quad (29)$$

We can obtain z by solving these three equations.

A straightforward calculation yields that these equations reduce to

$$4(a - b)^2 - (\sqrt{2(a + b) + 3} - 1)(\sqrt{2(a + b) + 3} + 3)^3 = 0, \quad (30)$$

where

$$a = \frac{1}{r^{\frac{1}{3}}} \left(\frac{r}{2} - \cos \theta + c_1 \sin^2 \theta \frac{1}{z} \right), \quad (31)$$

$$b = r^{\frac{1}{3}} \left(\frac{1}{2r} - \cos \theta + c_2 \sin^2 \theta \frac{1}{z} \right). \quad (32)$$

It can be checked that the above equations have a unique real positive solution of z . Eqs. (30)-(32) have been used for the analysis in this work.

In the special case of a “back-to-back” pair production, in which Eq. (14) holds, we recover the result obtained by taking the massless limit of the formula in Ref. [10],

$$(M_{T2})^2 \Big|_{\text{back-to-back}} = 2(|\mathbf{p}_T^{B,1}| |\mathbf{p}_T^{B,2}| + \mathbf{p}_T^{B,1} \cdot \mathbf{p}_T^{B,2}) = 2p_1 p_2 (1 + \cos \theta). \quad (33)$$

(ii) $m_X \neq 0$ **case:** Generalization of the above result for the case with massive X , i.e., $m_X \neq 0$, is straightforward. In this case, the M_{T2} variable is defined by Eq. (2) with $m_B = 0$. The same argument as above shows that Eq. (13) holds also in this case.

Calculating in the same way as above, it can be shown that

$$(M_{T2})^2 = m_X^2 + 2p_1 p_2 z \quad (34)$$

with z being the solution of Eq. (30) with

$$a = \frac{1}{r^{\frac{1}{3}}} \left(\frac{r}{2} - \cos \theta + c_1 \sin^2 \theta \frac{1}{z} - \frac{r \sin^2 \theta}{2} \frac{m_X^2}{p_2^2} \frac{1}{z^2} \right), \quad (35)$$

$$b = r^{\frac{1}{3}} \left(\frac{1}{2r} - \cos \theta + c_2 \sin^2 \theta \frac{1}{z} - \frac{\sin^2 \theta}{2r} \frac{m_X^2}{p_1^2} \frac{1}{z^2} \right). \quad (36)$$

For the case with massive X , this expression for M_{T2} is valid for any values of c_1 and c_2 . The existence of a unique positive real solution of z can also be checked.

References

- [1] H. Pagels and J. R. Primack, Phys. Rev. Lett. **48** (1982) 223;
M. Viel, J. Lesgourgues, M. G. Haehnelt, S. Matarrese and A. Riotto, Phys. Rev. D **71** (2005) 063534 [arXiv:astro-ph/0501562].
- [2] C. G. Lester and D. J. Summers, Phys. Lett. B **463** (1999) 99 [arXiv:hep-ph/9906349];
A. Barr, C. Lester and P. Stephens, J. Phys. G **29** (2003) 2343 [arXiv:hep-ph/0304226].
- [3] K. Kawagoe, T. Kobayashi, M. M. Nojiri and A. Ochi, Phys. Rev. D **69** (2004) 035003 [arXiv:hep-ph/0309031].
- [4] C. G. Lester and A. J. Barr, JHEP **0712** (2007) 102 [arXiv:0708.1028 [hep-ph]];
W. S. Cho, K. Choi, Y. G. Kim and C. B. Park, arXiv:0709.0288 [hep-ph];
A. J. Barr, B. Gripaios and C. G. Lester, JHEP **0802** (2008) 014 [arXiv:0711.4008 [hep-ph]];
W. S. Cho, K. Choi, Y. G. Kim and C. B. Park, JHEP **0802** (2008) 035 [arXiv:0711.4526 [hep-ph]].
M. M. Nojiri, Y. Shimizu, S. Okada and K. Kawagoe, arXiv:0802.2412 [hep-ph].

- [5] F. E. Paige, S. D. Protopopescu, H. Baer and X. Tata, [arXiv:hep-ph/0312045].
- [6] G. Marchesini, B. R. Webber, G. Abbiendi, I. G. Knowles, M. H. Seymour and L. Stanco, Comput. Phys. Commun. **67** (1992) 465;
 G. Corcella *et al.*, JHEP **0101** (2001) 010 [arXiv:hep-ph/0011363];
 G. Corcella *et al.*, [arXiv:hep-ph/0210213].
- [7] E. Richter-Was, [arXiv:hep-ph/0207355].
- [8] K. Hamaguchi, E. Nakamura, S. Shirai and T. T. Yanagida, arXiv:0804.3296 [hep-ph].
- [9] B. C. Allanach *et al.*, in *Proc. of the APS/DPF/DPB Summer Study on the Future of Particle Physics (Snowmass 2001)* ed. N. Graf, *In the Proceedings of APS / DPF / DPB Summer Study on the Future of Particle Physics (Snowmass 2001)*, Snowmass, Colorado, 30 Jun - 21 Jul 2001, pp P125 [arXiv:hep-ph/0202233].
- [10] C. G. Lester and A. J. Barr, in Ref. [4];
 W. S. Cho, K. Choi, Y. G. Kim and C. B. Park, JHEP **0802** (2008) 035, in Ref. [4].